



PC 752

# NEUTRACIDE 64

ONE-STEP DISINFECTANT, GERMICIDAL,  
DETERGENT AND DEODORANT

DISINFECTANT - PSEUDOMONACIDAL - STAPHYLOCIDAL - SALMONELLACIDAL - BACTERICIDAL  
FUNGICIDAL - MILDEWSTATIC - \*VIRUCIDAL - SANITIZER

**ACTIVE INGREDIENTS:**  
Didecyl dimethyl ammonium chloride ..... 2.54%  
n-Alkyl (C<sub>14</sub> 50%, C<sub>12</sub> 40%, C<sub>16</sub> 10%)  
dimethyl benzyl ammonium chloride ..... 1.69%  
**INERT INGREDIENTS:**..... 95.77%  
Total ..... 100.00%

EPA Reg. No. 47371-131-4313  
EPA Est. No. 4313-TX-1



**KEEP OUT OF REACH  
OF CHILDREN  
DANGER**  
See back panel for  
precautionary statements  
and First Aid.

ITEM#  
**752**



www.carrollco.com

A multi-purpose, neutral pH, germicidal detergent and deodorant effective in hard water up to 200 ppm (calculated as CaCO<sub>3</sub>) in the presence of a moderate amount of soil (5% organic serum) according to the AOAC Use-dilution Test. Disinfects, cleans, and deodorizes in one labor-saving step.

Kills 99.9% of Influenza A virus associated with the flu. Kills Pandemic 2009 H1N1 influenza A virus (formerly called swine flu).

For use in hospitals, nursing homes, schools, colleges, universities, commercial and industrial institutions, office buildings, veterinary clinics, animal life science laboratories, zoos, federally inspected meat and poultry plants, tack shops, pet shops, airports, kennels, hotels, motels, breeding establishments, and grooming establishments.

Disinfects, cleans, and deodorizes the following hard, nonporous, inanimate surfaces: floors, walls, non-medical metal surfaces, non-medical stainless steel surfaces, glazed porcelain, plastic surfaces (such as polypropylene, polystyrene, acrylic, etc.).

For use in: federally inspected meat and poultry plants on all hard, nonporous surfaces in inedible product processing areas, non-processing areas, and/or exterior areas. All surfaces must be thoroughly rinsed with potable water.

Effective against the following pathogens: **Bacteria:** Pseudomonas aeruginosa, Staphylococcus aureus, Salmonella enterica, Acinetobacter calcoaceticus, Bordetella bronchiseptica, Chlamydia psittaci, Community Associated Methicillin-Resistant Staphylococcus aureus - [CA-MRSA] (NRS384) (USA300), Community Associated Methicillin-Resistant Staphylococcus aureus - [CA-MRSA] (NRS123) (USA400), Enterobacter aerogenes, Enterobacter cloacae, Enterococcus faecalis-Vancomycin Resistant (VRE), Escherichia coli, Fusobacterium necrophorum, Klebsiella pneumoniae, Legionella pneumophila, Listeria monocytogenes, Pasteurella multocida, Proteus mirabilis, Proteus vulgaris, Salmonella enteritidis, Salmonella typhi, Salmonella typhimurium, Serratia marcescens, Shigella flexneri, Shigella sonnei, Staphylococcus aureus - Methicillin Resistant (MRSA), Staphylococcus aureus - Vancomycin Intermediate Resistant (VISA), Staphylococcus epidermidis, Streptococcus faecalis, Streptococcus pyogenes [Strep].

**Viruses:** Adenovirus type 4, \*Hepatitis B Virus (HBV), \*Hepatitis C Virus (HCV), \*Herpes Simplex Virus Type 1, \*Herpes Simplex Virus Type 2, \*Human coronavirus, \*HIV-1 (AIDS virus), \*Influenza A virus/Hong Kong, Influenza Virus\*\* [\*\*\*in association with the organism Influenza A/Hong Kong], \*Rotavirus [WA], \*Respiratory syncytial virus (RSV), \*Rubella virus, \*SARS Associated Coronavirus, \*Vaccinia, **Animal Viruses:** \*Avian Influenza H5N1, \*Avian polyomavirus, \*Canine distemper, \*Equine Herpes Virus Type 1, \*Feline coronavirus, \*Feline leukemia, \*Feline panleukopenia virus, \*Feline picornavirus, \*Infectious bovine rhinotracheitis, \*Infectious bronchitis (Avian IBV), \*Newcastle Disease, \*Porcine parvovirus, \*Pseudorabies (PRV), \*Rabies, \*Transmissible Gastroenteritis virus (TGE), \*Fungi: Aspergillus niger, Candida albicans, Trichophyton mentagrophytes (Athlete's Foot Fungus).

\*ATCC & antibiotic-resistant strain    \*\*antibiotic-resistant strain only

PELIGRO: SI NO PUEDE LEER EN INGLES, PREGUNTE A SU SUPERVISOR SOBRE LAS INSTRUCCIONES DE USO APROPIADAS ANTES DE TRABAJAR CON ESTE PRODUCTO.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

<b>DISINFECTION</b> DILUTION: 1:64 (660 PPM quat)	<b>SANITIZATION</b> DILUTION: 1:64 (660 PPM quat)	<b>2 OUNCES PER GALLON OF WATER</b>
---	---	---

This product is not for use on medical device surfaces.

**DISINFECTING/CLEANING/DEODORIZING DIRECTIONS:** Remove heavy soil deposits from surface, then thoroughly wet surface with a use-solution of 2 ounces of the concentrate per gallon of water. The use-solution can be applied with a cloth, mop, sponge, or coarse spray or by soaking. For sprayer applications, use a coarse spray device. Spray 6-8 inches from the surface, rub with a brush, cloth or sponge. Do not breathe spray. Let solution remain on surface for a minimum of 10 minutes. Rinse or allow to air dry. Rinsing of floors is not necessary unless they are to be waxed or polished. This product is not for use on food contact surfaces. **For Human Coronavirus**, treated surfaces must remain wet for 1 minute. Wipe dry with a clean cloth, sponge or mop or allow to air dry. **For Influenza Virus Type A**, treated surfaces must remain wet for 2 minutes. Wipe dry with a clean cloth, sponge or mop or allow to air dry.

Prepare a fresh use-solution daily or more often if the solution becomes visibly dirty or diluted.

**TOILET BOWLS:** Swab bowl with brush to remove heavy soil prior to cleaning or disinfecting. Clean by applying use-solution around the bowl and up under the rim. Stubborn stains may require brushing. To disinfect, first pre-clean to remove heavy soil, then remove or expel over the inner trap the residual bowl water. Pour in three ounces of the diluted solution. Swab the bowl completely using a scrub brush or mop, making sure to get under rim. Let stand for 10 minutes or overnight, then flush.

**Fungicidal Directions:** For use in areas such as locker rooms, dressing rooms, shower and bath areas and exercise facilities follow disinfection directions.

**MILDEWSTATIC INSTRUCTIONS:** Will effectively control the growth of mold and mildew, plus the odors caused by them when applied to hard, nonporous surfaces such as walls, floors, and table tops. Apply use-solution of 2 ounces per gallon of water with a cloth, mop, sponge, or coarse spray. Make sure to wet all surfaces completely. Let air dry. Repeat application weekly or when growth reappears.

**TO SANITIZE NON-FOOD CONTACT HARD NON-POROUS SURFACES:** Mix 2 oz. of product per gallon of water to sanitize hard nonporous surfaces. Treated surfaces must remain wet for 60 seconds. Wipe dry with a sponge, mop, or cloth or allow to air dry. Do not use in food preparation, food handling and food storage areas. This product is not to be used on glassware, dishware, cookware or utensils.

**To control the growth of mold and mildew on large inflatable non-porous plastic and rubber structures (animals, promotional items, moonwalks, obstacle course, play and exercise equipment):** Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use solution of 2 ounces per gallon of water with a cloth, mop, sponge, or coarse spray. Make sure to wet all surfaces completely. Let air dry. Repeat application weekly or when growth reappears. Ventilate buildings and enclosed spaces. Do not use equipment until treatment has been absorbed.

Manufactured for:



2900 W. Kingsley Rd., • Garland, Texas 75041 • 800.527.5722  
www.carrollclean.com

Medical Emergency 1-800-535-5053