



CLEAR LEMON

**DISINFECTANT, VIRUCIDE* AND CLEANER
FOR HOSPITAL, INSTITUTIONAL AND INDUSTRIAL USE**

*HUMAN IMMUNODEFICIENCY VIRUS, TYPE 1 (HIV-1)

ACTIVE INGREDIENT:

Alkyl (C14 50%, C12 40%, C16 10%)

dimethyl benzyl ammonium chloride 3.3%

INERT INGREDIENTS:..... 96.7%

**DILUTED
FOR USE**

**ITEM#
776**

The product in this container is diluted as directed on the pesticide product label

Follow the directions for use on the pesticide label when applying this product.

Si tiene preguntas, comuníquese con su supervisor.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. For use on precleaned, hard, inanimate surfaces. For heavy soil or organic matter, a precleaning step is required. Rinse with clear water. Apply disinfectant solution of two ounces per gallon of water with cloth or mop to toilets, restroom floors, shower stalls, diaper pails and garbage cans. Wet all surfaces thoroughly. Use four ounces per gallon of water for sick rooms, hospitals, or clinics. Keep surfaces wet for at least 10 minutes. Prepare a fresh solution daily or when use solution becomes visibly dirty.

PELIGRO: SI NO PUEDE LEER EN INGLES, PREGUNTE A SU SUPERVISOR SOBRE LAS INSTRUCCIONES DE USO APROPIADAS ANTES DE TRABAJAR CON ESTE PRODUCTO.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive. Keep Out of Reach of Children. Causes irreversible eye damage and skin burns. Harmful if swallowed. Do not get in eyes, on skin or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses), protective clothing and rubber gloves when handling. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. **IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

EPA Reg No. 4313-22

EPA Est. No. 4313-TX-1

1PS9900776_1

Manufactured for:



2900 W. Kingsley Rd., • Garland, Texas 75041

800.527.5722 • www.carrollclean.com

DANGER: KEEP OUT OF REACH OF CHILDREN